BP706T · B.Pharm Semester VII · Theory Notes

Pharmacy Practice

Unit IV — Patient-Oriented Services (9 hours)

Unit IV (9 hours)

Pills & Python Notes

From-zero teaching standard: every technical term is defined before use.

Syllabus coverage: Unit IV (9 hours).

PILLS & PYTHON

Pharmacy • Data • AI • Healthcare

Instagram @pillsandpython · pillsandpython.com · YouTube @PillsAndPython

CHAPTER 10: Medication Adherence and Non-Adherence

10.1 Learning Objectives

By the end of this chapter, the student will be able to:

1 Define medication adherence and non-adherence.

2 List and explain the factors that affect adherence.

3 Describe the pharmacist's role in improving medication adherence.

4 Describe tools used to monitor/assess adherence (pill counts, diaries, questionnaires — described generically).

5 Suggest strategies to overcome non-adherence.

6 Explain why adherence matters for treatment outcomes.

10.2 Definitions

Medication adherence: The extent to which a patient takes their medicines exactly as prescribed — the right drug, right dose, right time, right route, for the right duration.

Non-adherence (non-compliance): The extent to which a patient does NOT take medicines as prescribed — missing doses, taking wrong doses, taking at wrong times, or stopping treatment early.

Intentional non-adherence: The patient deliberately decides not to follow the prescription (e.g., stops a drug because of side effects or disbelief in its need).

Unintentional non-adherence: The patient fails to follow the prescription without meaning to (e.g., forgets doses, misunderstands instructions, cannot afford the medicine).

Persistence: Continuing to take a prescribed medicine over the full recommended duration without stopping early or having long gaps.

Adherence monitoring: The process of checking and measuring how well a patient is following the prescribed medicine regimen.

10.3 Terminology Table

Term	Meaning
Regimen	The complete prescribed plan: which medicines, what doses, at what times, for how long
Dose	The amount of medicine taken at one time
Compliance (older term)	An older word for adherence; "adherence" is preferred because it implies partnership, not just obedience
Concordance	An agreement between patient and prescriber about the treatment plan, reached through discussion

Term	Meaning
Pill count	An adherence check: counting tablets left in the patient's supply and comparing with what should be left
Medication diary	A daily log kept by the patient recording each dose taken (or missed)
Questionnaire (adherence)	A short set of standard questions asking the patient about missed doses and medicine-taking behaviour
Reminder system	Any aid that prompts the patient to take a dose on time (alarm, chart, family reminder)
Treatment outcome	The health result the therapy is meant to achieve

10.4 Simple Explanations

Concept 1: Adherence — taking medicines as agreed

- **Simple explanation:** Adherence means the patient follows the prescription faithfully: every dose, on time, for the full course. If the prescription says "one tablet twice daily for 10 days", adherence means 20 tablets taken correctly over 10 days.
- **Example:** A tuberculosis patient takes every single daily dose for the full months-long course — that is adherence; stopping after two months because "I feel better" is non-adherence.
- **Textual diagram:**

```
Prescription: 1 tablet, twice daily, 10 days (= 20 tablets)
        |
Patient takes all 20 correctly ──> ADHERENCE ──> best chance of cure
Patient misses several / stops early ──> NON-ADHERENCE ──> treatment may fail
```

- **Key takeaway:** Medicines only work if they are actually taken as directed.

Concept 2: Intentional vs unintentional non-adherence

- **Simple explanation:** Some patients *choose* not to take medicines (intentional — e.g., fear of side effects, belief the drug is unnecessary). Others *want* to take them but fail (unintentional — e.g., forgetfulness, confusion, cost). The two need different solutions: the first needs convincing and education; the second needs practical help.
- **Example:** Intentional — a patient stops blood-pressure tablets because "I feel fine now". Unintentional — an elderly patient misses the evening dose because she forgets.
- **Textual diagram:**

```
Non-adherence
     |
     +---> INTENTIONAL (decides not to take)
     |        Causes: side effects, disbelief, fear, stigma
     |        Fix: counselling, education, addressing beliefs
     |
     +---> UNINTENTIONAL (wants to, but fails)
              Causes: forgetfulness, complex regimen, cost, poor understanding
              Fix: reminders, simpler regimen, aids, affordability help
```

- **Key takeaway:** First find out *why* the patient is non-adherent — the reason decides the remedy.

Concept 3: Factors affecting adherence

- **Simple explanation:** Adherence is influenced by five groups of factors: the patient (beliefs, memory, age), the therapy (complexity, side effects, duration), the condition (no symptoms → less motivation; e.g., high blood pressure), the healthcare system (cost, distance, poor communication), and social support (family help or lack of it).
- **Example:** A patient with pain takes painkillers regularly (symptoms remind him), but the same patient skips his "silent" cholesterol tablet — the condition's nature affects adherence.
- **Key takeaway:** Non-adherence is rarely laziness — it has identifiable, fixable causes.

Concept 4: Pharmacist's role in improving adherence

- **Simple explanation:** The pharmacist is the medicine expert the patient meets most often. The pharmacist improves adherence by: explaining *why* each medicine matters, simplifying regimens where possible (with prescriber agreement), giving practical memory aids, checking understanding, following up, and spotting non-adherence early from refill patterns.
- **Example:** The pharmacist notices a patient refills a 30-day prescription every 45 days — a refill-gap clue to missed doses — and counsels the patient on the importance of daily dosing.
- **Key takeaway:** The pharmacist detects, understands, and fixes adherence problems at every patient contact.

Concept 5: Adherence monitoring tools — pill counts

- **Simple explanation:** Pill count: at a visit, the pharmacist counts how many tablets remain and compares with how many *should* remain if every dose was taken. Fewer tablets left than expected suggests doses were missed (or more taken than prescribed).
- **Example:** 30 tablets dispensed 15 days ago for twice-daily dosing → 0 should remain if fully adherent; 10 remaining suggests about 5 days of doses were missed.
- **Key takeaway:** Pill counts are simple and objective but can be fooled (patients may discard tablets).

Concept 6: Adherence monitoring tools — diaries

- **Simple explanation:** The patient keeps a simple daily diary — ticking each dose taken, noting missed doses and reasons. The pharmacist reviews it at each visit, spotting patterns (e.g., always missing the morning dose on workdays).
- **Example:** A diary shows the patient misses Sunday doses regularly — discussion reveals the patient sleeps in and skips breakfast-time tablets; the timing is shifted.
- **Key takeaway:** Diaries reveal *patterns* of missing doses, not just totals.

Concept 7: Adherence monitoring tools — questionnaires

- **Simple explanation:** A short set of standard, non-judgemental questions (e.g., "Do you ever forget to take your medicines?", "Have you ever stopped a medicine because you felt better?") asked in a friendly way. Honest answers estimate adherence level and uncover reasons.
- **Example:** A patient admits on a questionnaire to stopping tablets when feeling well — revealing intentional non-adherence that a pill count alone would not explain.
- **Key takeaway:** Questionnaires uncover the *reasons* behind non-adherence; ask without blaming.

Concept 8: Strategies to improve adherence

- **Simple explanation:** Strategies match the cause: educate and motivate (explain benefits, address fears), simplify (fewer tablets, suitable timings, combination products where appropriate and prescribed), remind (alarms, pill organizers, family support), follow up (phone calls, refill reminders), and reduce barriers (cost guidance, easy refill systems).
- **Example:** For a forgetful elderly patient: a weekly pill organizer + a morning alarm + a daughter who checks the organizer every Sunday — three simple strategies combined.
- **Key takeaway:** Combine education + simplification + reminders + follow-up; one strategy alone is rarely enough.

10.5 Classifications

Classification of non-adherence (by intent):

1 Intentional non-adherence — deliberate decision not to follow the regimen.

2 Unintentional non-adherence — failure despite willingness.

Classification of non-adherence (by pattern):

1 Missing single doses occasionally.

2 Taking drug holidays (skipping for days, then restarting).

3 Wrong timing or wrong dose.

4 Early discontinuation (stopping the course too soon).

5 Never starting (prescription filled but never taken).

Classification of factors affecting adherence (five groups):

1 Patient-related — beliefs, knowledge, memory, age, mental health, health literacy.

2 Therapy-related — complexity, number of tablets, side effects, taste, duration, frequent dosing.

3 Condition-related — silent diseases (no symptoms), chronic vs short-term illness, severity.

4 Healthcare-system-related — cost, distance to pharmacy, waiting time, poor communication, lack of follow-up.

5 Social/economic — family support, stigma, work schedule, cultural beliefs.

Classification of adherence monitoring tools:

1 Direct/objective — pill counts, refill records, diary review.

2 Indirect/subjective — patient questionnaires/interviews, clinical response assessment.

10.6 Mechanisms/Processes

Process: How non-adherence harms (the chain of consequences).

1 Doses are missed or stopped early.

2 Drug levels in the body fall below effective levels.

3 Disease is not controlled (blood pressure rises, infection returns, fits recur).

4 Complications develop; hospital admission may follow.

5 Stronger, costlier treatment becomes necessary; drug resistance can develop (e.g., in infections like tuberculosis).

Process: Pharmacist-led adherence improvement (step-by-step).

1 Detect — use refill gaps, pill counts, diaries, questionnaires, or clinical clues (poor control despite right prescription).

2 Understand — ask open, non-judgemental questions to find whether it is intentional or unintentional and which factors apply.

3 Plan — choose strategies matched to the cause; involve the patient in the plan (concordance).

4 Act — counsel, simplify aids, set reminders, coordinate with the prescriber if the regimen needs changing.

5 Follow up — re-check adherence at the next visit; praise improvement; adjust the plan.

Process: Using a questionnaire effectively.

1 Choose a private, unhurried moment.

2 Explain the purpose: "I ask everyone this so I can help you get the best from your medicines."

3 Ask short, simple, non-blaming questions one at a time.

4 Listen fully; note reasons, not just yes/no.

5 Summarize what you heard and agree on next steps with the patient.

10.7 Textual Diagrams/Flowcharts

Diagram 1 — The adherence improvement cycle:

```
DETECT (refill gaps, pill count, diary, questionnaire,
        poor clinical control)
            |
            v
UNDERSTAND (intentional or unintentional?
            which of the 5 factor groups?)
            |
            v
PLAN with patient (education + simplification
            + reminders + barrier removal)
            |
            v
ACT (counsel, aids, alarms, follow-up schedule)
            |
            v
FOLLOW UP (re-measure adherence; praise progress;
            adjust plan if still poor) ──> back to DETECT
```

Diagram 2 — Matching strategy to cause:

```
Cause found                    Best-fit strategy
──────────────────────────────────────────────────────────
Forgets doses          ──>     Alarms, pill organizer, routine-linking
Doesn't understand     ──>     Simple counselling + written instructions
Fears side effects     ──>     Explain, reassure, report/prescriber review
Feels "no need"        ──>     Explain silent diseases; show progress markers
Complex regimen        ──>     Simplify timing; discuss options with prescriber
Cost problem           ──>     Affordable alternatives; refill planning
No family support      ──>     Involve a family member; follow-up calls
```

10.8 Infographic Concept

INFOGRAPHIC CONCEPT: "The Adherence Iceberg" — an iceberg illustration: above the water, the visible tip labelled "Missed doses"; below the water, the much larger hidden mass divided into five labelled blocks — Patient beliefs, Complex therapy, Silent disease, Cost & access, Weak support. A pharmacist in a small boat circles the iceberg with a magnifying glass, with the caption: "Treat the cause below the surface, not just the missed dose above it." Colour theme: deep blue water, white ice, orange highlights.

10.9 Tables

Table 1: Intentional vs unintentional non-adherence

Feature	Intentional	Unintentional
Patient's attitude	Decides not to follow	Wants to follow but fails
Common causes	Side effects, disbelief, fear, stigma, feeling better	Forgetfulness, confusion, complexity, cost
Key solution	Education, addressing beliefs, shared decision-making	Reminders, simplification, practical aids
Detection clue	Admits stopping; good knowledge but poor intake	Wants to comply; confused about regimen

Table 2: Adherence monitoring tools compared

Tool	How it works	Strengths	Limitations
Pill count	Count tablets left vs expected	Simple, objective, quick	Can be manipulated; no reason revealed
Medication diary	Patient logs each dose daily	Shows patterns and reasons	Needs literacy and honesty; effort for patient
Questionnaire	Standard short questions, friendly interview	Uncovers reasons; cheap	Subjective; patients may over-report good behaviour
Refill records	Check refill intervals at pharmacy	Objective; no patient effort	Assumes dispensed = taken
Clinical response	Disease markers (e.g., BP, sugar)	Shows real outcome	Poor control has many causes besides adherence

Table 3: Five factor groups with examples

Factor group	Example causes
Patient-related	Poor understanding, forgetfulness, depression, low health literacy
Therapy-related	Many tablets, 3–4 times daily dosing, bad taste, side effects, long duration
Condition-related	No symptoms (hypertension), chronic lifelong treatment, stigma (e.g., mental illness)
Healthcare-system-related	High cost, far pharmacy, long waits, rushed consultations, no follow-up
Social/economic	No family support, busy work hours, cultural beliefs against medicines

Table 4: Strategies matched to factor groups

Factor group	Strategies
Patient-related	Simple counselling, teach-back, written instructions, involve family
Therapy-related	Simplify timing, pill organizer, discuss regimen options with prescriber
Condition-related	Explain silent diseases, show progress charts, set review dates
Healthcare-system-related	Refill reminders, easy refill process, cost counselling, follow-up calls
Social/economic	Family engagement, link doses to daily routine, community support

10.10 Examples

1 **Unintentional:** An elderly man misses his evening tablet because he falls asleep early — solved with a morning-shifted schedule approved by the prescriber.

2 **Intentional:** A young woman stops her thyroid tablet because she read online that it causes weight gain — solved by counselling with correct information and prescriber discussion.

3 **Pill count detection:** 12 tablets remain of a 30-tablet twice-daily supply after 15 days → about 6 days of doses missed.

4 **Diary pattern:** A diary shows every Monday dose missed — the patient travels for work on Mondays; a travel pill case solves it.

5 **Questionnaire revelation:** "I stop when I feel better" reveals intentional early stopping of an antibiotic course — counselled on completing the course.

6 **Refill gap:** A 30-day medicine refilled every 50 days signals ~20 missed days per cycle.

7 **Combined strategy:** For a busy mother: one pill organizer + phone alarm + linking doses to meals + a refill reminder call from the pharmacy.

10.11 Analogies

1 **Adherence is like watering a plant daily** — missing a few days may not kill it, but irregular watering keeps it weak; regular watering keeps it thriving.

2 **A prescription is like a railway ticket** — buying it (getting the medicine) is not enough; you must board the right train at the right time (take it correctly) to reach the destination (cure).

3 **Intentional vs unintentional non-adherence is like missing school on purpose vs missing the bus** — one needs persuasion, the other needs an alarm clock.

4 **The pill organizer is like a weekly tiffin box** — each compartment packed in advance, so you never wonder "did I take it today?"

5 **Refill records are like a bank statement** — they don't show what you *spent*, but the gaps tell a story the pharmacist can read.

10.12 Pharmacy Relevance

Adherence is where pharmacy practice meets real-world results: the best prescription fails if the patient doesn't take it. The community and clinical pharmacist — the most frequently seen healthcare professional — is ideally placed to detect non-adherence, understand its cause, and fix it. Every refill visit is an adherence check; every counselling moment is an adherence intervention. For the B.Pharm student, adherence work is the clearest proof that pharmacists improve health outcomes, not just supply products.

10.13 India Context

In India, long treatment courses (tuberculosis, diabetes, hypertension), large joint families with shared medicine-taking routines, cost pressures, and varying health literacy make adherence both a challenge and a major opportunity for pharmacists. National programmes for tuberculosis and other chronic diseases rely heavily on treatment supporters and follow-up — roles where pharmacists contribute through counselling, refill tracking and reminder systems. Simple, low-cost tools — pill organizers, diary cards in local language, family involvement, and refill reminder calls — work well in the Indian community-pharmacy setting, as per the latest programme guidelines/notifications (confirm the current edition).

10.14 Key Points

- Adherence = taking medicines exactly as prescribed; non-adherence = any deviation.
- Non-adherence is intentional (deliberate) or unintentional (despite willingness).

- Five factor groups: patient, therapy, condition, healthcare system, social/economic.
- Pharmacist's role: detect → understand → plan with patient → act → follow up.
- Tools: pill counts, medication diaries, questionnaires, refill records, clinical response — each with strengths and limits.
- Strategies must match the cause: educate, simplify, remind, follow up, remove barriers.
- Non-adherence causes treatment failure, complications, resistance, and higher costs.

10.15 Common Confusions

1 **Adherence vs compliance:** "Compliance" sounds like obedience; "adherence" implies the patient is a partner in the plan. Modern practice prefers "adherence" (and "concordance" for the agreement itself).

2 **Dispensed = taken:** A filled prescription does not mean the medicine was swallowed — refill records and pill counts only *suggest* adherence; they don't prove it.

3 **Feeling better = cured:** Many patients stop treatment when symptoms fade (e.g., antibiotics, blood-pressure drugs); the disease process often continues silently.

4 **One tool is enough:** No single tool measures adherence perfectly — combine tools (e.g., questionnaire + refill record) for a truer picture.

5 **Non-adherence = patient's fault:** Usually it is a system of causes (complex regimen, cost, poor communication) — blame fixes nothing; understanding fixes a lot.

10.16 Common Mistakes

1 Scolding or blaming the patient for missed doses — this closes communication.

2 Assuming the patient understood the regimen without checking (no teach-back).

3 Using only one adherence tool and treating its result as final.

4 Recommending a strategy that doesn't match the cause (e.g., an alarm for a patient who deliberately stopped the drug).

5 Ignoring refill-gap clues at the dispensing counter.

6 Forgetting to follow up after giving adherence advice.

7 Overlooking cost as a cause of non-adherence.

8 Counselling in a rushed, noisy setting where the patient can't ask questions.

10.17 Mnemonics

- **Adherence definition — "RDRRD":** **R**ight drug, **D**ose, **R**oute, **R**ight time, **D**uration.
- **Factor groups — "PTCHS":** **P**atient, **T**herapy, **C**ondition, **H**ealthcare system, **S**ocial/economic — "Patients Take Care; Health (system) Supports".
- **Pharmacist's role — "DUPAF":** **D**etect, **U**nderstand, **P**lan, **A**ct, **F**ollow up.

- **Tools — "PQDRC": P**ill count, **Q**uestionnaire, **D**iary, **R**efill records, **C**linical response.
- **Strategies — "ESRF": E**ducate, **S**implify, **R**emind, **F**ollow up.

10.18 Exam Summary

- Define adherence and non-adherence; distinguish intentional from unintentional with examples.
- Classify factors affecting adherence into five groups with examples.
- Describe the pharmacist's role in improving adherence (detect → understand → plan → act → follow up).
- Describe adherence monitoring tools: pill counts, medication diaries, questionnaires (generic descriptions), refill records, clinical response — with strengths and limitations.
- Suggest strategies to overcome non-adherence, matched to causes.
- Explain the consequences of non-adherence.

10.19 One-Minute Revision

Adherence = right drug, dose, route, time, duration. Non-adherence: intentional (chooses not to) vs unintentional (wants to but fails). Five factor groups: Patient, Therapy, Condition, Healthcare system, Social/economic. Pharmacist's cycle: Detect → Understand → Plan → Act → Follow up. Tools: pill count (count vs expected), diary (patterns), questionnaire (reasons, ask kindly), refill records (gaps), clinical response (outcome). Strategies matched to cause: Educate, Simplify, Remind, Follow up. No single tool is perfect — combine them. Never blame; understand and fix.

10.20 Questions

Short Answer Questions

1 Define medication adherence and non-adherence.

2 Distinguish between intentional and unintentional non-adherence with one example each.

3 List the five groups of factors affecting adherence.

4 What is a pill count? How is it interpreted?

5 What is a medication diary?

6 List four strategies to improve adherence.

7 What is persistence in the context of drug therapy?

8 Give two consequences of non-adherence.

Long Answer Questions

1 Define medication adherence and non-adherence. Classify non-adherence and describe the five groups of factors affecting adherence with examples.

2 Describe the pharmacist's role in improving medication adherence, covering detection, understanding causes, planning, action and follow-up.

3 Describe the tools used to monitor patient medication adherence (pill counts, diaries, questionnaires, refill records, clinical response), with their strengths and limitations.

4 Suggest strategies to overcome non-adherence, explaining how each strategy matches a specific cause. Describe the chain of consequences when adherence fails.

MCQs (with answer keys)

1 Medication adherence means:

a) Buying the medicine from any shop b) Taking medicines exactly as prescribed — right drug, dose, time, route, duration c) Taking medicines only when symptoms appear d) Sharing medicines with family members **Answer: b**

1 A patient stops tablets deliberately because of fear of side effects. This is:

a) Unintentional non-adherence b) Intentional non-adherence c) Good adherence d) Persistence **Answer: b**

1 An elderly patient misses doses because she forgets. This is:

a) Intentional non-adherence b) Unintentional non-adherence c) Concordance d) Overdosage **Answer: b**

1 In a pill count, 10 tablets remain when 0 should remain. This suggests:

a) Perfect adherence b) About 5 days of twice-daily doses were missed c) The patient took extra doses d) Nothing — pill counts are meaningless **Answer: b**

1 Which factor group includes "complex dosing schedule and bad taste"?

a) Patient-related b) Therapy-related c) Condition-related d) Social/economic **Answer: b**

1 The best first step when non-adherence is detected is to:

a) Scold the patient b) Ask open, non-judgemental questions to understand the cause c) Immediately change the prescription d) Ignore it **Answer: b**

1 A refill record showing a 30-day supply refilled every 50 days suggests:

a) Excellent adherence b) About 20 days of missed doses per cycle c) Over-adherence d) A dispensing error **Answer: b**

1 Which strategy best fits forgetfulness?

a) A louder scolding b) Alarms, pill organizer, linking doses to daily routine c) Stopping the medicine d) Doubling the next dose **Answer: b**

Viva Questions

1 Define adherence in one sentence.

2 How do you distinguish intentional from unintentional non-adherence in an interview?

3 What does a refill gap tell you?

4 Why is a questionnaire alone not enough to judge adherence?

5 Give one strategy each for forgetfulness, fear of side effects, and high cost.

6 What is "teach-back" in counselling, and why does it help adherence?

7 Why do patients with "silent" diseases like hypertension often show poor adherence?

Application-Based Questions

1 A diabetic patient's blood sugar remains high despite a correct prescription. At the pharmacy you find his 30-day tablets are refilled every 48 days. Walk through your detect → understand → plan → act → follow-up approach for this patient.

2 An elderly patient living alone takes six different medicines and often confuses them. Design a practical, low-cost adherence support plan using at least three strategies.

3 A young mother admits she stops her child's antibiotic once the fever settles. Explain how you would counsel her, addressing her specific belief without blaming her.

4 You want to start a simple adherence-monitoring service in your community pharmacy. Describe which tools you would use, how you would record findings, and how you would follow up patients.

10.21 References

- Parthasarathi, G., Hansen, K. N. and Nahata, M. C. A Textbook of Clinical Pharmacy Practice.
- Parmar, N. S. Health Education and Community Pharmacy.
- Hassan, W. E. Hospital Pharmacy.
- Merchant, S. H. and Qadry, J. S. A Textbook of Hospital Pharmacy.
- Harman, R. J. Handbook of Pharmacy – Health Care.
- Shargel, L. Comprehensive Pharmacy Review.

CHAPTER 11: Patient Counselling Techniques and Communication Skills

Inside This Chapter

- What patient counselling means and why pharmacists do it
- The numbered steps of a counselling session with a textual flowchart
- Counselling materials and aids (written, visual, digital)
- Communication skills every pharmacist needs
- Barriers to effective counselling and strategies to overcome them

- Pharmacy relevance, India context, mnemonics, and exam questions

Learning Objectives

After studying this chapter, you should be able to:

1 Define patient counselling and list its main objectives.

2 Describe each step of patient counselling in correct order.

3 Draw the counselling steps as a flowchart in the exam.

4 Name the types of counselling materials and aids, with one example of each.

5 List the verbal and non-verbal communication skills a pharmacist uses.

6 Identify barriers to effective counselling (patient, pharmacist, and environment related).

7 Suggest at least one strategy to overcome each type of barrier.

8 Explain how good counselling improves medicine adherence and patient safety.

Definitions

- **Patient counselling** — A one-to-one conversation in which the pharmacist gives the patient the information needed to use medicines safely and correctly, and checks that the patient understood.
- **Adherence** — Taking the medicine exactly as advised: right dose, right time, for the right duration.
- **Communication skill** — The ability to share information clearly and understand the other person — using words, voice, body language, and listening.
- **Verbal communication** — Sharing information through spoken or written words.
- **Non-verbal communication** — Sharing meaning without words: facial expressions, eye contact, gestures, posture.
- **Barrier** — Anything that blocks clear communication between the pharmacist and the patient.
- **Counselling aid** — Any tool (leaflet, chart, model, video) that helps explain the counselling message.
- **Feedback** — The pharmacist asking the patient to repeat or demonstrate, to confirm understanding.

Terminology Table

Term	What it means in simple words	How to remember
Counselling	Private talk to teach a patient how to use medicine correctly	Like coaching before a match
Adherence	Taking medicine exactly as told	"Adhere" = stick to the plan
Verbal	Using words (spoken or written)	Voice + words
Non-verbal	Body language — face, eyes, hands, posture	"Non" = without words
Barrier	Something that blocks the message	A wall between you and the patient

Term	What it means in simple words	How to remember
Aid	A helper tool that makes explaining easier	"Aid" = help
Feedback	Checking: "Please repeat what I said"	Two-way traffic, not one-way
Privacy	A private space where the patient feels safe to talk	"Private" = just for them
Empathy	Feeling with the patient, not pitying them	"Walk in their shoes"
Jargon	Technical words only professionals understand	Language that blocks, not builds

Concept-by-Concept Explanations

1. What Is Patient Counselling?

Concept. Patient counselling is a planned, private, face-to-face conversation in which the pharmacist explains everything the patient must know about their medicine — and then verifies that the patient can actually use it correctly.

Simple explanation. The doctor prescribes; the pharmacist teaches the patient *how to take it*. It is not just handing over the medicine. It is teaching — and then checking that the patient really learned.

Example. A patient gets an inhaler for asthma. The pharmacist explains what the inhaler is for, demonstrates the steps (shake → breathe out → press → breathe in slowly → hold breath), asks the patient to demonstrate it back, and confirms the patient can do it.

Textual flowchart.

```
Patient arrives with prescription
        |
        v
Pharmacist GREETS and creates privacy
        |
        v
Pharmacist EXPLAINS medicine clearly
        |
        v
Pharmacist DEMONSTRATES (e.g., inhaler use)
        |
        v
Patient REPEATS / DEMONSTRATES back
        |
        v
  Correct? ----NO----> Re-explain simply
        | YES
        v
Pharmacist CLOSES with summary + invitation to return
```

Key takeaway. Counselling = explain + demonstrate + verify understanding. If you only explained, you did not finish the job.

2. Objectives of Patient Counselling

Concept. Counselling has clear goals — it exists to make treatment work and keep the patient safe.

Simple explanation. Objectives (goals) of counselling:

1 Make sure the patient knows the medicine's name and purpose ("what is it for").

2 Teach correct dose, timing, and duration.

3 Teach how to take/store the medicine (with food? in the fridge?).

4 Warn about side effects and what to do if they appear.

5 Improve adherence — the patient actually finishes the full course.

6 Prevent medicine errors and harm.

7 Encourage the patient to ask questions and return for help.

Example. A mother collects antibiotics for her child. Objective: she must know the dose (5 mL twice daily), the duration (complete all 5 days even if the child feels better), storage (keep the reconstituted bottle in the refrigerator), and warning signs (return if the child gets a rash).

Key takeaway. Every counselling goal answers one patient question: What? How much? When? How long? What if something goes wrong? Whom do I ask?

3. Steps in Patient Counselling

Concept. Counselling follows fixed, ordered steps so nothing important is missed.

Simple explanation. The steps, in order:

1 **Greet and welcome the patient** — introduce yourself, be polite, learn the patient's name.

2 **Ensure privacy** — move to a quiet corner; the patient may be shy about a disease.

3 **Collect information** — ask about other medicines, allergies, pregnancy, diseases, habits (alcohol, smoking).

4 **Explain the prescription** — medicine name, purpose, dose, how and when to take it.

5 **Explain important warnings** — side effects, things to avoid (alcohol, driving), storage.

6 **Demonstrate when needed** — inhalers, injections (e.g., insulin pen), eye drops, syrup measuring.

7 **Verify understanding** — ask the patient to repeat key points or demonstrate; this is called *feedback*.

8 **Answer questions and close** — invite questions, summarize, give contact details for follow-up.

Example. Step 3 example: before dispensing a painkiller, the pharmacist asks, "Do you take any medicine for blood pressure?" — discovering a risk that changes the advice.

Textual diagram — the steps as a ladder.



Key takeaway. The steps are Greet → Privacy → Ask → Explain → Warn → Demonstrate → Verify → Close. Step 7 (verify) is what separates real counselling from a lecture.

4. Counselling Materials and Aids

Concept. Some patients forget spoken words quickly. Aids — written, visual, or digital tools — support the spoken message.

Simple explanation. Types of counselling materials:

- **Written materials** — patient information leaflets, reminder cards, medicine schedule charts, labels with pictograms (pictures).
- **Visual aids** — posters, flip charts, models of the body (e.g., a heart model), demonstration devices (placebo inhalers), pill organizers showing days of the week.
- **Digital aids** — short videos on a tablet, reminder alarms on the patient's phone, apps that track doses.
- **Language-matched aids** — materials written in the patient's own language, not only English.

Example. For an elderly patient on five medicines, the pharmacist gives a weekly pill organizer (Monday–Sunday boxes) plus a simple chart in the local language with a picture of each tablet and when to take it.

Key takeaway. Spoken words fade; a leaflet stays on the fridge. Always pair talking with something the patient can take home or keep.

5. Communication Skills for Pharmacists

Concept. Communication skill = how you deliver the message and receive the patient's concerns. It has spoken, unspoken, and listening parts.

Simple explanation.

- **Verbal skills:** speak slowly, use simple everyday words (avoid medical jargon), keep sentences short, give information in small chunks.
- **Non-verbal skills:** maintain gentle eye contact, smile, nod while listening, face the patient, avoid crossing arms (looks closed-off).

- **Listening skills:** let the patient finish speaking, repeat back to confirm ("So you are saying..."), ask open questions ("What worries you about this medicine?") instead of only yes/no questions.
- **With prescribers:** communicate problems politely and professionally — e.g., "Doctor, this patient seems allergic to penicillin; may I suggest discussing an alternative?" Use written notes or calls, not public criticism.

Example. Bad: "Take this stat and monitor glycemic indices." Good: "Take this tablet right now, before you leave. It lowers your blood sugar — check your sugar again tomorrow morning."

Key takeaway. Talk *with* the patient, not *at* them. The patient's understanding is the pharmacist's responsibility.

6. Barriers to Effective Counselling and How to Overcome Them

Concept. A barrier is anything that stops the message from landing. Barriers come from three sides: the patient, the pharmacist, and the environment.

Simple explanation.

Source	Barrier	Strategy to overcome
Patient	Cannot read well (low literacy)	Use pictures, demonstrations, simple words
Patient	Language difference	Use the patient's language; interpreter or translated leaflets
Patient	Shyness, fear, or distrust	Be warm, ensure privacy, never judge
Patient	Memory problems (elderly)	Give written notes, pill organizer, family involvement
Patient	Beliefs ("medicines are harmful")	Listen respectfully, explain benefits and risks honestly
Pharmacist	Using jargon	Replace technical terms with everyday words
Pharmacist	Rushing, no time	Keep counselling short but structured; prioritize key points
Pharmacist	Poor listening (interrupting)	Wait, nod, confirm before replying
Environment	Noisy, crowded counter	Create a small counselling corner, even a marked space
Environment	No privacy	Counsel in a quieter area; speak softly

Example. A patient who cannot read gets a leaflet — useless. The pharmacist instead shows pictures on a chart and demonstrates with the actual medicine bottle. Barrier identified, strategy applied.

Key takeaway. Every barrier has an antidote. A good pharmacist first notices the barrier, then adjusts the method.

Classifications

Classification of counselling materials:

```
Counselling aids
├── Written: leaflets, schedule charts, reminder cards, pictogram labels
├── Visual: posters, flip charts, body models, placebo devices, pill organizers
├── Digital: videos, phone alarms, dose-tracking apps
└── Language-matched: any of the above in the patient's own language
```

Classification of communication skills:

```
Communication skills
├── Verbal: simple words, short sentences, small chunks, no jargon
├── Non-verbal: eye contact, smile, open posture, nodding
├── Listening: let them finish, repeat back, open questions
└── Professional: polite prescriber communication, written follow-up
```

Classification of barriers:

```
Barriers
├── Patient-side: illiteracy, language, shyness, memory, beliefs
├── Pharmacist-side: jargon, rushing, poor listening, judgemental tone
└── Environment-side: noise, crowding, no privacy, interruptions
```

Mechanisms / Processes

The counselling process (input → process → output):

```
INPUT (before counselling)
  prescription + patient history + medicine knowledge
        |
        v
PROCESS (the 8 steps)
  greet → privacy → ask → explain → warn → demonstrate → verify → close
        |
        v
OUTPUT (after counselling)
  informed patient + correct use + better adherence + fewer errors
```

The feedback loop mechanism. Feedback (step 7) is a loop, not a line:

```
Explain → Patient repeats → Correct?
      YES → move on
      NO  → explain differently → patient repeats again → Correct? → ...
```

This loop continues until the patient gets it right. Never skip it because the patient "looks smart."

Textual Diagrams and Flowcharts

Diagram 1 — Full counselling flow (exam-ready):



Diagram 2 — Barrier → strategy map:

BARRIER		STRATEGY
Patient can't read	→	Pictures + demonstration
Wrong language	→	Own-language leaflet / interpreter
Shy patient	→	Privacy + warm, non-judging tone
Forgets easily	→	Written notes + pill organizer + family
Pharmacist jargon	→	Simple everyday words
Rushing	→	Prioritize: dose, timing, warnings first
Noisy counter	→	Counselling corner, speak clearly
No privacy	→	Quiet spot, soft voice

INFOGRAPHIC CONCEPT: A one-page poster titled "The 8 Steps of Patient Counselling" showing a staircase of 8 numbered steps (Greet → Privacy → Ask → Explain → Warn → Demonstrate → Verify → Close) with a small icon beside each (smiley face, curtain, question mark, medicine bottle, warning triangle, hand demonstrating, speech bubble with tick, waving hand). At the bottom, a ribbon reads: "Explain + Demonstrate + VERIFY — counselling is complete only when the patient shows they understood." Side panel: "Top 5 barriers" as five brick-wall icons, each cracked open by a strategy key.

Tables

Table 1 — Jargon vs simple words (what to say instead):

Medical jargon (avoid)	Simple words (use)
Take this stat	Take this right now
Twice daily	Two times a day — morning and night
On an empty stomach	30 minutes before food
Hypersensitivity	Allergy / reaction
Adverse effect	Side effect / unwanted effect
Discontinue	Stop taking
Store in a cool place	Keep away from heat and sunlight

Table 2 — Which aid for which situation:

Situation	Best aid
Patient cannot read	Pictures, demonstrations, pill organizer
Complex device (inhaler, insulin pen)	Placebo device for hands-on practice
Many medicines	Weekly pill organizer + picture chart
Patient speaks another language	Translated leaflet or interpreter
Patient forgets doses	Phone alarm, reminder card, family support

Examples

1. **Inhaler counselling:** A new asthma patient receives an inhaler. The pharmacist demonstrates with a placebo device, then asks the patient to show the steps back. The patient forgets to shake it — the pharmacist re-demonstrates. Correct technique achieved on the second try.
2. **Antibiotic course:** A father collects antibiotics for his son. The pharmacist stresses: "Give it for all 5 days, even if he feels well on day 3. Stopping early lets the infection return stronger." Gives a written schedule card.
3. **Elderly polypharmacy:** A 70-year-old takes six medicines. The pharmacist prepares a weekly pill organizer and a chart with tablet pictures and timings in the local language, and explains it to both the patient and the daughter.
4. **Prescriber communication:** The pharmacist notices a patient is prescribed two medicines with the same ingredient under different brand names. The pharmacist phones the doctor politely: "Doctor, both brands contain the same drug — the patient may take a double dose by mistake. Would you like to keep one?" The doctor agrees. A double-dose error is prevented.
5. **Side-effect warning:** A patient starts iron tablets. The pharmacist warns: "Your stools may turn dark — that is normal and harmless. Take it on an empty stomach, but if it upsets your stomach, take it after food." The patient is not frightened when the change appears.

Analogies

- **Counselling is like teaching someone to drive, not just handing them car keys.** The prescription is the car; the patient still needs lessons (and a test) before driving alone.

- **Feedback is like a cricket umpire's signal.** You don't assume the ball crossed the boundary — you check the signal. Don't assume understanding — check it.
- **A barrier is like static on a phone call.** The message is fine, but noise blocks it. Remove the noise (or change the channel — pictures instead of words), and the message lands.
- **Counselling aids are like a recipe book.** A cooking teacher can explain a dish, but the recipe card on the fridge is what the cook follows at home.

Pharmacy Relevance

Patient counselling is one of the most visible clinical services a pharmacist provides. It directly:

- Improves adherence, so treatments actually work.
- Prevents medication errors (wrong dose, wrong technique, dangerous combinations).
- Builds public trust in pharmacists as healthcare providers, not just medicine sellers.
- Reduces hospital visits caused by preventable medicine problems.
- Is a core part of pharmaceutical care and medication therapy management.

India Context

- In India, the pharmacist at a retail/community pharmacy is often the most accessible health professional — many patients visit the pharmacy before (or instead of) a doctor, especially in rural and semi-urban areas.
- Language diversity means counselling must often happen in the patient's mother tongue; Hindi/regional-language leaflets and pictogram-based labels are especially valuable.
- Low health literacy in some populations makes verbal explanation + demonstration + pictures more effective than written English leaflets alone.
- Pharmacists in India counsel on a wide range of products: prescription medicines, OTC products, and traditional/herbal preparations — the same structured steps apply.
- Pharmacy regulations in India (as per the latest guidelines/notifications — confirm the current edition) define the pharmacist's responsibilities toward patients; counselling is part of professional duty.

Key Points

1. Counselling = private teaching + verification of understanding.
2. Objectives: correct use, correct dose/timing/duration, warnings, adherence, error prevention.
3. Eight steps in order: Greet → Privacy → Ask → Explain → Warn → Demonstrate → Verify → Close.
4. Step 7 (verify — patient repeats/demonstrates) is the step that makes counselling real.
5. Aids: written (leaflets, charts), visual (models, placebo devices, pill organizers), digital (videos, alarms), language-matched.

6 Communication skills: simple verbal language, open body language, active listening, polite prescriber communication.

7 Barriers: patient-side (literacy, language, shyness, memory, beliefs), pharmacist-side (jargon, rushing, poor listening), environment-side (noise, no privacy).

8 Every barrier has a strategy — match the strategy to the barrier.

Common Confusions

- **Counselling vs dispensing:** Dispensing = giving the medicine. Counselling = teaching about the medicine. Both happen at the counter, but they are different jobs.
- **Counselling vs advising:** Advice can be casual ("drink water"). Counselling is structured, private, and verified.
- **Verify vs ask "Do you understand?":** "Do you understand?" gets a polite "yes" even when the patient understood nothing. Verification means the patient *shows* (repeats or demonstrates) — that is the only real proof.
- **Empathy vs sympathy:** Sympathy = feeling sorry *for* someone ("poor you"). Empathy = understanding *with* them ("this must be difficult for you") and responding helpfully.

Common Mistakes

1 **Skipping verification** — the single most common failure; assuming understanding instead of checking it.

2 **Using jargon** — saying "take this stat" or "twice daily" to a patient who needs "right now" and "morning and night."

3 **Counselling in a crowd** — discussing a sensitive disease loudly at a busy counter, so the patient stays silent and asks nothing.

4 **Not asking about allergies or other medicines** — missing information that could prevent harm.

5 **No take-home aid** — the patient forgets everything by the time they reach home.

6 **Arguing with the patient** — contradicting the patient's beliefs instead of listening and gently correcting.

Mnemonics

- **Steps of counselling: "GREAT PVC"** — **G**reet, **P**rivacy, ask (**R**eview history — collect **E**ssential info)... A cleaner version: **"Good Pharmacists Always Explain Warnings, Demonstrate, Verify, Close"** = **G**reet, **P**rivacy, **A**sk, **E**xplain, **W**arn, **D**emonstrate, **V**erify, **C**lose.
- **Barriers: "PLE"** — **P**atient-side, pharmacist-side (**L**istener-side — the pharmacist), **E**nvironment-side.
- **Communication: "VNL"** — **V**erbal (words), **N**on-verbal (body), **L**istening (ears).

Exam Summary

Patient counselling is a structured, private conversation in which the pharmacist teaches the patient to use medicines safely and verifies understanding. Its objectives are correct medicine use, adherence, side-effect awareness, and error prevention. The eight steps are: (1) greet, (2) ensure privacy, (3) collect patient information (allergies, other medicines), (4) explain the prescription, (5) warn about side effects and storage, (6) demonstrate devices, (7) verify understanding through patient repeat-back, and (8) close with summary and follow-up. Counselling aids include written materials (leaflets, charts), visual aids (models, placebo devices, pill organizers), digital aids (videos, alarms), and language-matched materials. Communication skills cover verbal (simple words), non-verbal (eye contact, open posture), listening (open questions, no interrupting), and professional communication with prescribers. Barriers arise from the patient (literacy, language, shyness, memory, beliefs), the pharmacist (jargon, rushing, poor listening), and the environment (noise, no privacy) — each matched with a strategy such as pictures, interpreters, privacy, written notes, and plain language.

One-Minute Revision

Counselling = teach + verify. 8 steps: Greet, Privacy, Ask, Explain, Warn, Demonstrate, Verify, Close — remember "Good Pharmacists Always Explain Warnings, Demonstrate, Verify, Close." Never skip Verify. Aids: leaflets, charts, models, placebo devices, pill organizers, videos, own-language materials. Communicate simply, listen fully, watch body language. Barriers: patient (literacy/language/shyness), pharmacist (jargon/rushing), environment (noise/no privacy) — match each with a strategy.

Questions

Short Answer (2–3 marks)

1 Define patient counselling in one or two sentences.

2 List any four objectives of patient counselling.

3 Name the eight steps of patient counselling in order.

4 Give two examples each of written and visual counselling aids.

5 What is "feedback" in counselling, and why is it essential?

6 Name two patient-side and two environment-side barriers to counselling.

7 Convert to simple words: "Take this stat, twice daily, on an empty stomach."

8 Why is privacy important during patient counselling?

Long Answer (10 marks)

1 Describe the steps involved in patient counselling with a flowchart. Explain why the verification step is the most important.

2 Discuss the types of educational materials used in patient counselling. Give suitable examples for an illiterate patient and an elderly patient on multiple medicines.

3 Classify the barriers to effective counselling. For each category, suggest practical strategies a community pharmacist can use to overcome them.

4 "Communication is the heart of patient counselling." Discuss the verbal, non-verbal, and listening skills a pharmacist needs, with examples.

MCQs (with answer key)

1 Which step of counselling involves the patient repeating or demonstrating the instructions?

a) Greet b) Explain c) Verify d) Close — **Answer: c**

1 A pill organizer is an example of:

a) Verbal aid b) Visual aid c) Digital aid d) Prescriber aid — **Answer: b**

1 Which is a patient-side barrier to counselling?

a) Noisy counter b) Jargon use c) Low literacy d) Rushing — **Answer: c**

1 The best response to a shy patient is:

a) Speak loudly b) Ensure privacy and a warm tone c) Give a long leaflet d) Skip counselling — **Answer: b**

1 "Do you understand?" is a poor check because:

a) It takes too long b) Patients may say yes without understanding c) It is rude d) It needs writing — **Answer: b**

Viva Questions

1 What is the difference between dispensing and counselling?

2 Name the eight counselling steps without looking.

3 Why should you ask about allergies before dispensing?

4 What would you do if a patient cannot read your leaflet?

5 How do you communicate a prescription problem to a doctor politely?

Application-Based Questions

1 A 65-year-old man with poor eyesight collects three new medicines. Plan a complete counselling session: steps, aids, and barrier strategies.

2 A young mother is worried because her child's antibiotic syrup must be "shaken well and refrigerated." She seems confused. How would you counsel her, step by step?

3 You notice a prescription contains two brands of the same drug. Write the exact polite message you would give the prescriber, and explain why this communication matters.

References

- Hassan, W. E. *Hospital Pharmacy*. Lea & Febiger. (Preferably latest edition)
- Harman, R. J. *Handbook of Pharmacy – Health Care*. Pharmaceutical Press. (Preferably latest edition)
- Merchant, S. H. and Qadry, J. S. *A Textbook of Hospital Pharmacy*. CBS Publishers & Distributors. (Preferably latest edition)

- Parmar, N. S. *Health Education and Community Pharmacy*. CBS Publishers & Distributors. (Preferably latest edition)
- Parthasarathi, G., Hansen, K. N. and Nahata, M. C. *A Textbook of Clinical Pharmacy Practice*. Universities Press. (Preferably latest edition)
- Shargel, L. *Comprehensive Pharmacy Review*. Lippincott Williams & Wilkins. (Preferably latest edition)

CHAPTER 12: Health Screening Services and Telemedicine

Inside This Chapter

- What health screening means and why it matters
- Four screening services pharmacists can provide: blood pressure, blood sugar, BMI, lung function
- Screening vs diagnosis — and when to refer the patient
- Telemedicine: definition, how it works, the pharmacist's role
- Advantages and limitations of telemedicine
- Pharmacy relevance, India context, mnemonics, and exam questions

Learning Objectives

After studying this chapter, you should be able to:

1 Define health screening and explain its importance.

2 Describe the four screening methods: blood pressure measurement, blood sugar monitoring, BMI calculation, and lung function testing.

3 Explain the difference between screening and diagnosis.

4 State the pharmacist's role in health screening services.

5 List the signs that require referral to a doctor.

6 Define telemedicine and explain how a telemedicine consultation works.

7 Describe the pharmacist's role in telemedicine.

8 Compare the advantages and limitations of telemedicine.

Definitions

- **Health screening** — A quick, simple check done on people who feel healthy (or at risk), to find early warning signs of disease *before* symptoms appear.

- **Diagnosis** — The doctor's formal decision about exactly which disease a patient has, after full examination and tests. Screening is NOT diagnosis.
- **Referral** — Sending the patient to a doctor or specialist because the screening result needs expert review.
- **Blood pressure (BP)** — The force of blood pushing against the walls of the arteries (blood vessels).
- **Blood sugar (blood glucose)** — The amount of sugar (glucose) present in the blood.
- **BMI (Body Mass Index)** — A number calculated from height and weight that tells whether a person's weight is in a healthy range.
- **Lung function test** — A simple breathing test (for example, peak flow) that measures how well air moves in and out of the lungs.
- **Telemedicine** — Medical care given from a distance using phones, video calls, or the internet, instead of the patient visiting the clinic in person.
- **Teleconsultation** — A consultation (meeting with a doctor) done through telemedicine — e.g., a video call.

Terminology Table

Term	What it means in simple words	How to remember
Screening	Early check for warning signs	Like checking smoke before fire
Diagnosis	Doctor's final name for the disease	Only a doctor "names" it
Referral	Sending the patient to the right expert	"Refer" = pass to a specialist
Blood pressure	Force of blood on vessel walls	Water pressure in a pipe
Blood glucose	Sugar level in the blood	Fuel gauge for the body
BMI	Height–weight number for healthy weight	A weight report card
Peak flow	How fast you can blow air out	Blowing out birthday candles — speed measured
Telemedicine	Doctor visit through a screen	"Tele" = far, medicine from far
Teleconsultation	The actual video/phone appointment	A clinic visit without travel
Follow-up	Checking the patient again later	"Follow" = keep watching

Concept-by-Concept Explanations

1. Health Screening: Concept and Importance

Concept. Screening looks for the *early, silent* stages of disease — before the patient feels sick — so that treatment can start early, when it works best.

Simple explanation. Many diseases (high blood pressure, diabetes) grow quietly for years without symptoms. A screening test is a quick check that raises a flag: "Something looks unusual — see a doctor." Screening does not name the disease; it only says "needs a closer look."

Example. A 45-year-old man feels perfectly fine. At the pharmacy, a BP check shows a repeatedly high reading. He is referred to a doctor, who confirms hypertension early. Because it was caught early, lifestyle changes and one medicine control it — before any heart damage.

Textual diagram — screening vs diagnosis:

```
SCREENING (pharmacist)              DIAGNOSIS (doctor)
──────────────────────              ──────────────────
Quick, simple test        --->      Detailed examination + tests
"Something looks unusual" --->      "You HAVE hypertension"
Raises a flag             --->      Names the disease, starts treatment
Done on healthy/at-risk   --->      Done on the flagged patient
people
```

Key takeaway. Screening finds suspects; diagnosis convicts. The pharmacist screens and refers — the doctor diagnoses and treats.

2. Blood Pressure Measurement (Screening)

Concept. The pharmacist measures the patient's BP with a digital BP monitor (or manual cuff) and compares the reading with normal ranges.

Simple explanation. Steps: the patient sits quietly and rests for a few minutes; the cuff is wrapped on the upper arm; the machine shows two numbers (upper = systolic, lower = diastolic). One high reading is not enough — repeat on another day or advise the patient to recheck before concluding anything.

Example. A woman's reading is high on one visit. The pharmacist does not declare "you have hypertension." Instead: "Your reading is on the higher side today. Please come back in two days, or see your doctor for a proper check."

Key takeaway. Pharmacists *measure and flag* BP; they never *diagnose* hypertension.

3. Blood Sugar Monitoring (Screening)

Concept. Using a small glucometer, the pharmacist checks the patient's blood sugar from a finger-prick drop of blood.

Simple explanation. Steps: clean hands, prick the fingertip with a lancet, place the drop on the test strip, read the number on the meter. Fasting (empty stomach, morning) and random readings mean different things — the pharmacist notes *when* the test was done. A high reading = refer to the doctor, not "you have diabetes."

Example. A man who drinks a lot of water and urinates frequently gets a random sugar check at the pharmacy — the reading is high. The pharmacist refers him to a doctor with the recorded reading, date, and time. The doctor later confirms diabetes.

Key takeaway. Record three things with every sugar reading: the number, the date/time, and whether the patient had eaten. Context makes the number meaningful.

4. BMI Calculation (Screening)

Concept. BMI = weight (kg) ÷ height (m) ÷ height (m). The resulting number places the person in a weight category (underweight / healthy / overweight / obese ranges, as per the latest guidelines — confirm the current edition).

Simple explanation. Steps: measure weight on a scale, measure height, calculate BMI, compare with the standard ranges. BMI is a *screening* tool — it suggests who may need lifestyle advice or a doctor's review;

it is not a full health verdict.

Example. A patient weighs 80 kg and is 1.65 m tall. BMI = 80 ÷ 1.65 ÷ 1.65 ≈ 29.4 — in the overweight range. The pharmacist gives simple lifestyle counselling (balanced diet, daily walk) and suggests a doctor's review.

Key takeaway. BMI is a two-minute calculation that starts a useful conversation about weight and health.

5. Lung Function Testing (Screening)

Concept. A peak flow meter is a small handheld device: the patient blows hard into it, and it shows how fast air leaves the lungs — a rough measure of airway openness.

Simple explanation. Steps: the patient stands, takes a deep breath, seals lips around the mouthpiece, and blows out as hard and fast as possible. The best of three tries is recorded. Low readings in a person with cough/wheezing = refer to the doctor; pharmacists do not diagnose asthma or COPD.

Example. A young man with a long-standing night cough blows a low peak flow. The pharmacist records the reading and refers him to a doctor, who investigates further.

Key takeaway. The peak flow meter is a screening whistle — it signals "airways may be narrow, get checked," nothing more.

6. When to Refer

Concept. Referral means recognizing the limit of screening and handing the patient to the right expert.

Simple explanation. Refer to a doctor when:

- Any screening reading is repeatedly abnormal (high BP, high sugar, low peak flow).
- The patient has symptoms (chest pain, breathlessness, fainting, severe headache).
- Readings are dangerously extreme on a single check.
- The patient is pregnant, a child, or elderly with multiple diseases — extra caution.
- The patient asks for diagnosis or treatment — that is the doctor's job.

Example. A patient's BP reads very high AND he complains of a severe headache. The pharmacist does not wait — this is an urgent referral: "Please go to the hospital/doctor right away."

Textual flowchart — screen-and-refer:



Key takeaway. Screen → record → decide: normal (reassure), abnormal (refer soon), emergency (refer now). Always give the patient a written record of readings to carry to the doctor.

7. Telemedicine: Definition and How It Works

Concept. Telemedicine = healthcare delivered across distance using communication technology — the doctor and patient meet over video/phone instead of in the same room.

Simple explanation. How a teleconsultation works, step by step:

1 The patient books an appointment (phone/app/helpline).

2 At the appointment time, a video or audio call connects patient and doctor.

3 The doctor asks questions, sees the patient on video, reviews reports the patient shares.

4 The doctor gives advice and sends an electronic prescription.

5 The pharmacist can then dispense against that prescription and counsel the patient (in person or by phone).

6 Follow-up visits happen the same way.

Example. A patient in a remote village video-calls a doctor in the city for a skin rash. The doctor prescribes an ointment electronically; the local pharmacist dispenses it and explains how to apply it — the patient never travelled.

Key takeaway. Telemedicine moves the *consultation* to the patient, instead of moving the patient to the consultation.

8. Pharmacist's Role in Telemedicine

Concept. The pharmacist is the local, physical link in a telemedicine chain.

Simple explanation. What pharmacists do:

- Help patients connect (set up the call, especially for elderly patients).

- Share screening records (BP/sugar readings) with the doctor during the call.
- Dispense medicines against electronic prescriptions received after teleconsultation.
- Counsel the patient on the newly prescribed medicines (Chapter 11 skills apply).
- Follow up by phone: "Are you taking the medicine? Any problems?"

Example. An elderly woman cannot operate a video app. The pharmacist sets up the call at the pharmacy, stays nearby, shows the doctor her BP diary, and after the call explains the new prescription to her in her own language.

Key takeaway. In telemedicine, the doctor may be far — the pharmacist is near. The pharmacist makes the distant care actually work for the patient.

9. Advantages and Limitations of Telemedicine

Concept. Telemedicine is a powerful tool with clear boundaries.

Simple explanation.

Advantages:

- Saves travel time and cost, especially for rural patients.
- Faster access to specialists who may be far away.
- Useful for follow-up visits and minor/chronic conditions.
- Keeps care going during emergencies or when travel is difficult.
- Digital records make follow-up easier.

Limitations:

- No physical examination — the doctor cannot touch, listen to the chest, or feel the abdomen.
- Needs a good phone/internet connection and basic digital skills — not everyone has these.
- Not suitable for emergencies (chest pain, accidents, severe illness).
- Privacy and data-security concerns with video calls and digital prescriptions.
- Some patients (elderly, very ill) still need in-person care.

Example. Telemedicine works well for a diabetes follow-up ("show me your sugar diary"). It does NOT work for a road-accident injury — that patient needs a hospital immediately.

Key takeaway. Telemedicine is excellent for follow-ups, minor problems, and distant access — but it can never fully replace physical examination or emergency care.

Classifications

Classification of screening services in this chapter:

```
Pharmacist-led screening
├── Blood pressure measurement (digital monitor / cuff)
├── Blood sugar monitoring (glucometer, finger-prick)
├── BMI calculation (weight + height → number)
└── Lung function testing (peak flow meter)
        ALL are screening, NONE is diagnosis
```

Classification of referral urgency:

```
Referral
├── Routine: abnormal but stable → see doctor soon, carry written readings
└── Urgent: emergency symptoms/extreme readings → hospital/doctor NOW
```

Classification of telemedicine roles of the pharmacist:

```
Pharmacist in telemedicine
├── Before: help patient connect, prepare screening records
├── During: assist the call, share records with the doctor
└── After: dispense e-prescription, counsel, phone follow-up
```

Mechanisms / Processes

The screening process (mechanism):

```
OFFER screening (walk-in / camp / routine visit)
        |
        v
PREPARE patient (rest before BP, clean hands before sugar test)
        |
        v
PERFORM test correctly (right technique, right device)
        |
        v
RECORD result (number + date + time + context, e.g., fasting or not)
        |
        v
INTERPRET as screening only ("higher than usual" — never a disease name)
        |
        v
DECIDE: reassure / routine referral / urgent referral
        |
        v
DOCUMENT and give the patient a written copy
```

The teleconsultation process:

```
Book appointment → Video/audio call connects → Doctor questions +
patient shows reports → Doctor advises → E-prescription sent →
Pharmacist dispenses + counsels → Phone follow-up
```

Textual Diagrams and Flowcharts

Diagram 1 — Screening services at a glance:



Diagram 2 — Telemedicine flow:



INFOGRAPHIC CONCEPT: A one-page poster titled "Screen, Don't Diagnose" split into four quadrants — BP cuff, glucometer, weighing scale with height chart, peak flow meter — each with 3 tiny step icons and the rule "Record: number + date + time + context." Across the bottom, a traffic-light strip: GREEN (normal → reassure + healthy habits), YELLOW (abnormal → routine referral with written record), RED (emergency symptoms → urgent referral NOW). Side panel: "Telemedicine in 6 steps" as a phone-to-doctor-to-pharmacist chain, with a stethoscope-crossed-out icon labeled "No physical exam — know the limits."

Tables

Table 1 — The four screening services compared:

Service	Device/tool	What the patient does	What the pharmacist records	Key caution
Blood pressure	Digital BP monitor/cuff	Sits quietly, rests a few minutes	Two numbers, date, time	One high reading ≠ hypertension; recheck
Blood sugar	Glucometer, lancet, strip	Finger-prick drop of blood	Number, date, time, fasting or not	Note meal status — it changes meaning
BMI	Weighing scale + height measure	Stands on scale	Weight, height, calculated BMI	Screening tool only, not a full verdict
Lung function	Peak flow meter	Blows hard into mouthpiece	Best of 3 readings	Low reading → refer; never diagnose asthma

Table 2 — Screening vs diagnosis:

Point	Screening (pharmacist)	Diagnosis (doctor)
Purpose	Find early warning signs	Name the exact disease
Done on	Healthy or at-risk people	The flagged/symptomatic patient
Result says	"Needs a closer look"	"You have X; treatment is Y"
Equipment	Simple, quick tools	Full examination + lab tests

Table 3 — Telemedicine: advantages vs limitations:

Advantages	Limitations
No travel — saves time and money	No physical examination possible
Specialist access for remote areas	Needs internet + digital skills
Good for follow-ups and chronic care	Not for emergencies
Care continues when travel is hard	Privacy/data-security concerns
Digital records aid follow-up	Some patients still need in-person visits

Examples

1. **BP camp:** During a pharmacy health camp, a pharmacist screens 40 adults. Three have repeatedly high readings — each gets a written BP card and a routine referral. One has chest discomfort — urgent referral the same day.
2. **Sugar screening:** A 50-year-old with increased thirst gets a fasting sugar check at the pharmacy. The reading is high; the pharmacist records number, date, time, "fasting," and refers to the doctor. No disease name is mentioned.
3. **BMI counselling:** A college student calculates BMI in the overweight range at the pharmacy. The pharmacist gives simple diet and activity advice and suggests a doctor's review — no crash-diet promises.
4. **Peak flow:** A teenager with night-time coughing blows a low peak flow. The pharmacist records the best-of-three reading and refers to a doctor for evaluation.
5. **Telemedicine follow-up:** After a video consultation for hypertension, the doctor sends an e-prescription. The pharmacist dispenses it, counsels the patient on timing and side effects, and phones after a week to check adherence.

Analogies

- **Screening is like a smoke detector.** It doesn't tell you what is burning or how to put out the fire — it just beeps early so experts can act. Diagnosis is the firefighter naming and fighting the fire.
- **BMI is like a school report card's first page.** It gives a quick overall signal but doesn't tell the full story of the student.
- **Telemedicine is like a video call with family.** You can talk, see faces, and share news — but you can't hug them. Some things still need being there in person.
- **Referral is like passing the baton in a relay race.** The pharmacist runs the first lap (screening); the doctor runs the next (diagnosis and treatment). Dropping the baton — failing to refer — loses the race.

Pharmacy Relevance

- Health screening is a natural extension of community pharmacy: the pharmacist is accessible without an appointment.
- Screening builds the pharmacist's role as a preventive-care provider and strengthens public trust.
- Every screening is also a counselling opportunity (lifestyle advice, adherence, Chapter 11 skills).
- In telemedicine, the pharmacist is the indispensable local partner — connecting patients, dispensing e-prescriptions, and following up.
- Screening records kept by the pharmacist help doctors make faster, better decisions.

India Context

- In India, pharmacies are often the first (and sometimes only) health contact in rural and remote areas — screening at the pharmacy can catch silent diseases like hypertension and diabetes early in populations with limited doctor access.
- Government health programs run screening camps (for example, for diabetes and hypertension) in which pharmacists can participate; pharmacists should follow the latest guidelines/notifications for such programs — confirm the current edition.
- Telemedicine has grown rapidly in India, including government-supported teleconsultation services; pharmacists support it by helping digitally inexperienced patients connect and by dispensing and counselling on e-prescriptions.
- Language and digital-literacy barriers mean the pharmacist's hands-on help (setting up calls, explaining in the local language) is often what makes telemedicine actually usable for elderly and rural patients.
- Low-cost devices (digital BP monitors, glucometers) are widely available in Indian pharmacies, making these four screenings practical at the community level.

Key Points

1. Screening = early warning check on healthy/at-risk people; diagnosis = the doctor naming the disease.
2. Four pharmacist screening services: BP measurement, blood sugar monitoring, BMI calculation, lung function (peak flow) testing.
3. Every reading must be recorded with number + date + time + context (e.g., fasting or not).
4. Pharmacists screen and refer — they never diagnose.
5. Referral: routine (abnormal but stable → doctor soon) vs urgent (emergency signs → NOW).
6. Telemedicine = healthcare at a distance via phone/video/internet.
7. Teleconsultation flow: book → call → doctor advises → e-prescription → pharmacist dispenses + counsels → follow-up.
8. Pharmacist's telemedicine role: connect, share records, dispense, counsel, follow up by phone.

9 Advantages: access, no travel, follow-ups; Limitations: no physical exam, needs connectivity/skills, not for emergencies.

Common Confusions

- **Screening vs diagnosis:** A high BP reading at the pharmacy is NOT a diagnosis of hypertension. Only a doctor, after proper evaluation, diagnoses.
- **One abnormal reading vs a pattern:** A single high reading can come from stress, coffee, or a faulty technique. Repeat before worrying — but still record and advise recheck.
- **Telemedicine vs a phone chat:** A proper teleconsultation is a scheduled clinical appointment with records and a prescription — not casual medical advice over a call.
- **BMI vs fitness:** BMI is a screening number, not a full health certificate. A muscular athlete can have a high BMI and be healthy; the number starts a conversation, not ends it.

Common Mistakes

1 **Diagnosing from a screening test** — telling a patient "you have diabetes" after one glucometer reading. (Correct: record, explain it needs a doctor's check, refer.)

2 **Not recording context** — writing a sugar number without noting fasting/random, date, and time, making the number useless to the doctor.

3 **Skipping rechecks** — acting on a single BP reading taken when the patient rushed in.

4 **Delaying urgent referral** — advising "come back next week" when the patient has emergency symptoms.

5 **Using telemedicine for emergencies** — trying a video call for chest pain or an accident instead of sending the patient to hospital immediately.

6 **No written record for the doctor** — referring verbally without giving the patient their readings on paper.

Mnemonics

- **Four screenings: "BBBL"** — **B**P, **B**lood sugar, **B**MI, **L**ung function ("Big Brothers Bring Lungs").
- **Record every reading: "NDTC"** — **N**umber, **D**ate, **T**ime, **C**ontext (fasting or not).
- **Referral decision: "RUN"** — **R**outine (stable abnormal → soon), **U**rgent (emergency → NOW), **N**ormal (reassure).
- **Telemedicine flow: "C-CDEF"** — **C**all connects, **C**onsultation, **D**octor advises, **E**-prescription, **F**ollow-up (pharmacist dispenses + counsels in between).

Exam Summary

Health screening is a quick check on healthy or at-risk people to catch early warning signs before symptoms appear; it is not diagnosis, which is the doctor's job. Pharmacists can provide four screening

services: blood pressure measurement (cuff, patient rested, two numbers), blood sugar monitoring (glucometer finger-prick; record fasting or random), BMI calculation (weight ÷ height^2, compared with standard ranges), and lung function testing (peak flow meter, best of three). Every result is recorded with number, date, time, and context. Abnormal results lead to referral — routine (see doctor soon with written record) or urgent (emergency symptoms → hospital now). Telemedicine delivers healthcare at a distance through phone/video/internet: the patient books, a call connects patient and doctor, the doctor advises and sends an e-prescription, and the pharmacist dispenses, counsels, and follows up by phone. The pharmacist's role includes helping patients connect, sharing screening records, dispensing e-prescriptions, and counselling. Advantages include access without travel and specialist reach; limitations include no physical examination, dependence on connectivity and digital skills, unsuitability for emergencies, and privacy concerns.

One-Minute Revision

Screening finds warning signs; diagnosis names the disease — pharmacists screen, doctors diagnose. Four screenings: BP, blood sugar, BMI, peak flow — record Number, Date, Time, Context. Refer: Routine (soon) or Urgent (NOW) — "RUN". Telemedicine = care at a distance; pharmacist connects, shares records, dispenses e-prescriptions, counsels, follows up. Good for follow-ups and remote access; never for emergencies or physical exams.

Questions

Short Answer (2–3 marks)

1 Define health screening in one or two sentences.

2 How is screening different from diagnosis? (Two points)

3 List the four screening services a pharmacist can provide.

4 What four details must be recorded with every screening reading?

5 Define telemedicine.

6 Give two situations in which urgent referral is needed.

7 State two advantages and two limitations of telemedicine.

8 What is the pharmacist's role after a teleconsultation ends?

Long Answer (10 marks)

1 Describe the four health screening services a pharmacist can provide, including the method and one key caution for each. Explain when the pharmacist must refer the patient, with a flowchart.

2 Define telemedicine and explain the complete teleconsultation process step by step. Discuss the pharmacist's role at each stage.

3 "Screening is not diagnosis." Discuss this statement with reference to BP, blood sugar, BMI, and lung function screening, including the pharmacist's responsibilities and limits.

MCQs (with answer key)

1 Which of the following can a pharmacist do based on a screening test?

a) Diagnose hypertension b) Record and refer c) Start treatment d) Order surgery — **Answer: b**

1 BMI is calculated as:

a) height ÷ weight b) weight (kg) ÷ height (m) ÷ height (m) c) weight + height d) weight × height — **Answer: b**

1 A patient has severe headache with a very high BP reading. The pharmacist should:

a) Ask him to return next week b) Start a medicine c) Arrange urgent referral now d) Repeat the test tomorrow — **Answer: c**

1 Which is a limitation of telemedicine?

a) Saves travel time b) No physical examination possible c) Good for follow-ups d) Digital records — **Answer: b**

1 "Number, date, time, context" refers to:

a) Prescription format b) What to record with each screening reading c) Referral letter format d) Telemedicine steps — **Answer: b**

Viva Questions

1 What is the difference between screening and diagnosis?

2 Name the four screening services in this chapter.

3 Why must you note whether a sugar test was fasting or random?

4 When would you choose urgent referral over routine referral?

5 Can telemedicine replace a physical examination? Why or why not?

Application-Based Questions

1 At a pharmacy health camp, a 48-year-old man's BP reads high on a single check. He feels fine. Describe exactly what you would do, step by step, and what you would NOT do.

2 An elderly patient in a village needs a teleconsultation but cannot operate a smartphone. Plan the pharmacist's role before, during, and after the call.

3 A patient's glucometer reading at your pharmacy is high. Write the written record you would give him to carry to the doctor, and explain your referral advice.

References

- Hassan, W. E. *Hospital Pharmacy*. Lea & Febiger. (Preferably latest edition)
- Harman, R. J. *Handbook of Pharmacy – Health Care*. Pharmaceutical Press. (Preferably latest edition)
- Merchant, S. H. and Qadry, J. S. *A Textbook of Hospital Pharmacy*. CBS Publishers & Distributors. (Preferably latest edition)

- Parmar, N. S. *Health Education and Community Pharmacy*. CBS Publishers & Distributors. (Preferably latest edition)
- Parthasarathi, G., Hansen, K. N. and Nahata, M. C. *A Textbook of Clinical Pharmacy Practice*. Universities Press. (Preferably latest edition)
- Shargel, L. *Comprehensive Pharmacy Review*. Lippincott Williams & Wilkins. (Preferably latest edition)